arXiv:hep-ph/9711378v1 18 Nov 1997

University of Wisconsin - Madison

MADPH-97-1022
hep-ph/9711378
November 1997

On the LEP Measurements of the Rising Photon-Photon Total Cross Section

M. M. Block
Department of Physics, Northwestern University
Evanston, IL 60208

E. M. Gregores and F. Halzen
Department of Physics, University of Wisconsin
Madison, WI 53706

G. Pancheri
INFN - Laboratori Nazionali di Frascati
Frascati, Italy

Abstract

Using vector meson dominance we calculate the $\gamma\gamma$ total cross section from data on pp, $p\bar{p}$ and γp total cross sections. Our result agrees with recent high energy measurements performed by the L3 experiment at the LEP collider, not with the conflicting preliminary data obtained by OPAL.

Typeset using REVTEX

Two experiments at LEP have measured the high energy $\gamma\gamma$ total cross section [1]. While these measurements yield new information on its high energy behavior at center-of-mass energies in excess of $\sqrt{s} = 15$ GeV, they may represent the last opportunity to measure the $\gamma\gamma$ cross section, and the two data sets unfortunately disagree. We here point out that a simultaneous analysis of pp, $p\bar{p}$, γp and $\gamma\gamma$ total cross sections in the framework of vector dominance (VMD) nicely accommodates the L3 measurements. The analysis is sufficiently restrictive to exclude the preliminary OPAL results.

We have performed a simultaneous analysis of all data on forward scattering amplitudes involving hadrons and photons in the context of a model where high energy cross sections rise as a consequence of the increasing numbers of soft quarks and gluons populating the colliding particles [2]. This idea is implemented in an eikonal formalism where the total cross section is given by

$$\sigma_{\rm tot}(s) = 2P_{\rm had}\int \left[1 - e^{-\chi_{\rm I}}(s,b)\cos(\chi_{\rm R}(s,b)\right]\, d^2\vec{b}\,, \tag{1}$$

with the eikonal χ given by $\chi(s,b) = \chi_{\rm R}(s,b) + i\chi_{\rm I}(s,b)$. Here b is the impact parameter and $P_{\rm had}$ the probability that the particle interacts as a hadron. Its value is obviously one for the proton and antiproton, and of order α when the particle is a photon. Its value can be obtained from VMD. We will use the value $P_{\rm had} = 1/240$ for γp collisions and, therefore, $P_{\rm had} = (1/240)^2$ for $\gamma\gamma$ collisions.

The eikonal χ receives contributions from quark-quark, quark-gluon and gluon-gluon interactions, therefore

$$\begin{aligned}\chi(s,b) &= \chi_{qq}(s,b) + \chi_{qg}(s,b) + \chi_{gg}(s,b)\\ &= A(\mu_{qq},b)\sigma_{qq}(s) + A(\sqrt{\mu_{qq}\mu_{gg}},b)\sigma_{qg}(s) + A(\mu_{gg},b)\sigma_{gg}(s)\,,\end{aligned} \tag{2}$$

where $A(\mu,b)$ is the overlap function in impact parameter space and σ_{ij} the cross sections of the colliding partons. We simply parametrize $A(\mu,b)$ as the Fourier transform of a dipole form factor

$$A(\mu,b) = \frac{\mu^2}{96\pi}(\mu b)^3 K_3(\mu b)\,, \tag{3}$$

where $K_3(x)$ is the modified Bessel function of second kind. Since the "size" of quarks and gluons in the proton can be different, we obtain $\mu_{qq} = 0.89$ and $\mu_{gg} = 0.73$. This model where hadrons asymptotically evolve into black disks of partons nicely accommodates all data on pp and $p\bar{p}$ total cross sections as well as measurements of the differential elastic cross section and of the real-to-imaginary part of the forward scattering amplitude [3]. The results shown in Fig. 1 have been updated for a new measurement of the $p\bar{p}$ total cross section at the Fermilab collider [4].

Following reference [5] we obtain γp and $\gamma\gamma$ total cross sections from the assumption that, in the spirit of VMD, the photon is a 2 quark state in contrast with the proton which is a 3 quark state. The γp total cross section is obtained from the even eikonal (used in the pp and $p\bar{p}$ fit of Fig. 1) by the substitutions:

$$\begin{aligned}\sigma_{ij} &\rightarrow \tfrac{2}{3}\,\sigma_{ij}\\ \mu_i &\rightarrow \sqrt{\tfrac{3}{2}}\,\mu_i\,.\end{aligned}$$

To obtain the $\gamma\gamma$ total cross section, we now substitute

$$\begin{aligned} \sigma_{ij} &\to \tfrac{4}{9}\,\sigma_{ij} \\ \mu_i &\to \tfrac{3}{2}\,\mu_i \end{aligned}$$

into the nucleon-nucleon even eikonal.

The results are shown in Figs. 2, 3. The agreement with data is impressive. The formalism also accommodates the HERA measurements of elastic ρ photoproduction. It is clear that VMD selects the L3 and rejects the preliminary OPAL results [1].

We have performed this analysis using the model of reference [2], and repeated it by performing a calculation of the parton cross section in the mini-jet model [6]. The conclusions are identical and the fits indistinguishable. Details will be published elsewhere [3].

ACKNOWLEDGMENTS

One of us (MMB) was partially supported by Department of Energy grant DA-AC02-76-Er02289 Task B. This research was also supported in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation, by the U.S. Department of Energy under Grant No. DE-FG02-95ER40896, and by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP).

REFERENCES

[1] PLUTO Collaboration, Ch. Berger *et al.*, Phys. Lett. **B149**, 421 (1984); TPC/2γ Collaboration, H. Aihara *et al.*, Phys. Rev. D**41**, 2667 (1990); MD-1 Collaboration, S. E. Baru *et al.*, Z. Phys. **C53**, 219 (1992); L3 Collaboration, M. Acciarri *et al.*, Phys. Lett. **B408**, 450 (1997); F. Wäckerle, *"Total Hadronic Cross-Section for Photon-Photon Interactions at LEP"*, to be published in the Proceedings of the XXVII International Symposium on Multiparticle Dynamics, Frascati, September 1997, and Nucl. Phys. **B**, Proc. Suppl.

[2] M. M. Block, R. Fletcher, F. Halzen, B. Margolis and P. Valin, Phys. Rev. D**41**, 978 (1990).

[3] M. M. Block *et al.*, to be published.

[4] J. Orear, to be published in the Proceedings of the International Conference (7^{th} Blois Workshop) on Elastic and Diffractive Scattering - Recent Advances in Hadron Physics, Seoul, Korea, June 1997.

[5] R. S. Fletcher, T. K. Gaisser and F. Halzen, Phys. Rev. D**45**, 377 (1992); Phys. Rev. D**45**, 3279 (1992) (E); Phys. Lett. **B298**, 442 (1993).

[6] D. Cline, F. Halzen and J. Luthe, Phys. Rev. Lett. **31**, 491 (1973); P. l'Hereux, B. Margolis and P. Vallin, Phys. Rev. D**32**, 1681 (1985); L. Durand and H. Pi, Phys. Rev. Lett. **58**, 303 (1987); Phys. Rev. D**40**, 1436 (1989); V. Innocente, A. Capella and J. T. T. Van, Phys. Lett. **B213**, 81 (1988); B. Margolis *et al.*, Phys. Lett. **B213**, 221 (1988); B. Z. Kopeliovich, N. N. Nikolaev and I. K. Potashnikova, Phys. Rev. D**39**, 769 (1989); J. C. Collins and G. A. Ladinsky, Phys. Rev. D**43**, 2847 (1991).



FIG. 1. The total cross section for pp and $p\bar{p}$ scattering in mb, σ vs $\sqrt{s}$, the center-of-mass energy in GeV. The solid line and squares are for pp scattering and the dotted line and circles are for $p\bar{p}$ scattering.



FIG. 2. The total γp cross section in mb, $\sigma(\gamma p)$ vs $\sqrt{s}$, the center-of-mass energy in GeV.



FIG. 3. The total $\gamma\gamma$ cross section in nb, $\sigma(\gamma\gamma)$ *vs* $\sqrt{s}$, the center-of-mass energy in GeV.